



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

with O , must be circular; therefore, the central ellipsoid must be one of revolution. The last is a *non-sequitur*, because, as we have seen, if O is on the normal to a circular section, it satisfies the condition that the two parallel sections normal to d , one with centre at O and the other with centre at g , the centre of mass of the body, shall both be circular. Poinso't's reasoning shows no more than that O must be on a normal to a circular section of the central ellipsoid; it does not show that this circular section must be a principal section, although he gives that as his conclusion.

It must be confessed that, while it was shown above that under certain conditions there are four points for which the momental ellipsoid has two circular sections at right angles to each other, the analysis does not show conclusively that there are *only* four such points. We have discovered all such points as are situated on normals to central circular sections, but are there no such points elsewhere?

ON THE MASS OF TITAN.

By PROF. ORMOND STONE, University of Virginia, Va.

There is a numerical error in the value of a_6 as given on page 169, Vol. III of this journal. The correct value is approximately $-\frac{1}{2}a_3^2$. The introduction of this in equation (19), p. 166, adds other appreciable terms, the largest of which is $-3a_3a_6 = \frac{3}{2}a_3^3$. If, also, we take into account the term $m'P_0$ in ν , we may, in the place of equation (31), write

$$(2a_3P_0 + P_3)m' = \left[1 - 9 \left(\frac{n' - n}{n}\right)^2\right] a_3,$$

approximately; whence

$$m' = \frac{1}{4617}.$$

This change in the mass leads, of course, to corresponding changes in the values of the coefficients a_1 , n_1 , etc.